



From the currency rate quotations onto strings and brane world scenarios

D. Horváth, R. Pincak*

SORS Research a.s, 040 01 Kosice, Slovak Republic

ARTICLE INFO

Article history:

Received 4 May 2011

Received in revised form 29 May 2012

Available online 12 June 2012

Keywords:

Econophysics

Financial markets

String theory

ABSTRACT

In the paper, we study the projections of the real exchange rate dynamics onto the string-like topology. Our approach is inspired by the contemporary movements in the string theory. The string map of data is defined here by the boundary conditions, characteristic length, real valued and the method of redistribution of information. As a practical matter, this map represents the detrending and data standardization procedure. We introduced maps onto 1-end-point and 2-end-point open strings that satisfy the Dirichlet and Neumann boundary conditions. The questions of the choice of extra-dimensions, symmetries, duality and ways to the partial compactification are discussed. Subsequently, we pass to higher dimensional and more complex objects. The 2D-Brane was suggested which incorporated bid-ask spreads. Polarization by the spread was considered which admitted analyzing arbitrage opportunities on the market where transaction costs are taken into account. The model of the rotating string which naturally yields calculation of angular momentum is suitable for tracking of several currency pairs. The systematic way which allows one suggest more structured maps suitable for a simultaneous study of several currency pairs was analyzed by means of the Gâteaux generalized differential calculus. The effect of the string and brane maps on test data was studied by comparing their mean statistical characteristics. The study revealed notable differences between topologies. We review the dependence on the characteristic string length, mean fluctuations and properties of the intra-string statistics. The study explores the coupling of the string amplitude and volatility. The possible utilizations of the string theory approach in financial markets are slight.

© 2012 Elsevier B.V. All rights reserved.

1. Introduction

We are currently in the process of transfer of modern physical ideas into the neighboring field called econophysics. The physical statistical view point has proved fruitful, namely, in the description of systems where many-body effects dominate. However, standard, accepted by physicists, bottom-up approaches are cumbersome or outright impossible to follow the behavior of complex economic systems, where autonomous models encounter the intrinsic variability.

Digital economy is founded on data. In the paper, we suggest and analyze statistical properties of heuristics based on the currency rate data which are arranged to mimic the topology of the basic variants of the physical strings and branes. Our primary motivation comes from the actual physical concepts [1,2]; however, our realization differs from the original attempts in various significant details. The second aspect of our method is that it enables a transformation into a format which is useful for an analysis of a partial trend or relative fluctuations on the time scale window of interest.

* Corresponding author. Tel.: +421 907936147.

E-mail addresses: pincak@saske.sk, pincak@sors.com (R. Pincak).

As with most science problems, the representation of data is the key to efficient and effective solutions. The underlying link between our approach and the string theory may be seen in the switching from a local to a non-local form of the data description. This line passes from the single price to the multivalued collection of prices from the temporal neighborhood which we term here the string map. As we will see later, an important role in our considerations is played by the distance measure of the string maps. The idea of exploring the relationship between more intuitive geometric methods and financial data is not new. The discipline called the geometric data analysis includes many diverse examples of the conceptual schemes and theories grounded on the geometric representation and properties of data. Among them we can emphasize the tree network topology that exhibits usefulness in the studies of the world-trade network [3] and other network structures of the market constructed by means of inter-asset correlations [4,5]. The multivariate statistical method called *cluster analysis* deals with data mapping onto representative subsets called *clusters* [6]. Here we work on the concept that is based on projection data into higher dimensional vectors in the sense of the work [7,8]. Also, arguments based on the metrics are consistent with our efforts but not too obvious points in common with the original objectives of the nonlinear analysis.

The string theory development over the past 25 years achieved a high degree of popularity among physicists [9,10]. The reason lies in its inherent ability to unify theories that come from diverse physical spheres. The prime instrument of the unification represents the concept of extra dimension. The side-product of theoretical efforts can be seen in the elimination of the ultraviolet divergences of Feynman diagrams. However, despite the considerable achievements, there is a lack of the experimental verification of the original string theory. In contrast, in the present work we exploit time-series which can build the family of the string motivated models of boundary-respecting maps. In a narrow sense, the purpose of the present data-driven study is to develop statistical techniques for the analysis of these objects.

The work is organized as follows: In Section 2, we specify the data selection and data pretreatment. In Section 3, we introduce the notion of the string map of data time series. The symmetry of the maps is discussed in Section 4. To examine generality and specificity of these ideas, the calculations have been performed for several representative currencies and ask (buyer initiated) or bid (seller initiated) prices. In Section 5, we give a generalization for a higher dimensional case which encodes the ask-bid spread difference. In this section, we also discuss the problem of partial compactification (Section 5.1). The statistical picture (in Sections 6–8) is of primary interest. The analysis of fluctuations measured in the middle of a string by means of the statistical moments is carried out in Section 6.2. In Section 7, we focus on the statistics of intra-string degrees of freedom. The efforts are in particular justified by the analytical studies in Sections 7.1 and 7.2. They include a decomposition of internal string states into Fourier modes in Section 7.3. In Section 8, we will see how the arbitrage opportunities, including spread, may be characterized by means of the model of *polarized string*. In the same section, we briefly discuss a reformulation of the popular concept of the correlation sum in terms of the strings and branes. There are also indications of a synergy between the string amplitudes and volatility in Section 6.3. Efforts have been made to study inter-currency relations by means of the projections onto *rotating strings* (Section 9) and via a generalization of the derivative (Section 10).

2. Data analysis

First of all we need to mention some facts about data streams we analyzed. We analyze tick by tick data of EUR/USD, GBP/USD, USD/JPY, USD/CAD, USD/CHF major currency pairs from the OANDA market maker. We focused on the three month period within three selected periods of 2009 which capture moments of the financial crisis. The streams are collected in such a way that each stream begins with Monday. More precisely, we selected periods denoted as Aug.–Sep. (from Aug. 3 to Sep. 7), Sep.–Oct. (Sep. 7–Oct. 5) and Oct.–Nov. (Oct. 5–Nov. 2). At first, the data sample has been decimated – only each 10th tick was considered. This delimits results to the scales larger than 10 ticks. The mean time corresponding to the string length l_s in ticks is given by

$$T(l_s) = \langle t(\tau + l_s) - t(\tau) \rangle \simeq \frac{1}{\tau_{\text{up}} - \tau_{\text{dn}}} \sum_{\tau=\tau_{\text{dn}}}^{\tau_{\text{up}}} [t(\tau + l_s) - t(\tau)]. \quad (1)$$

Data for study of rotating strings and angular moments (see Section 9) were preformatted in a different way. In this case, the currency information has been projected onto the grid of the regularly spaced 10 s intervals.

3. One-dimensional maps

By applying standard methodologies of detrending one may suggest to converting original series of the quotations of the mean currency exchange rate $p(\tau)$ onto a series of returns defined by

$$\frac{p(\tau + h) - p(\tau)}{p(\tau + h)}, \quad (2)$$

where h denotes a tick lag between currency quotes $p(\tau)$ and $p(\tau + h)$, τ is the index of the quote. The mean $p(\tau) = (p_{\text{ask}}(\tau) + p_{\text{bid}}(\tau))/2$ is calculated from $p_{\text{ask}}(\tau)$ and $p_{\text{bid}}(\tau)$.

In the spirit of the string theory it would be better to start with the 1-end-point open string map

$$P^{(1)}(\tau, h) = \frac{p(\tau + h) - p(\tau)}{p(\tau + h)}, \quad h \in \langle 0, l_s \rangle \quad (3)$$

where superscript (1) refers to the number of endpoints.

Later, we may use the notation $P\{p\}$ which emphasizes the functional dependence upon the currency exchange rate $\{p\}$. It should also be noted that the use of P highlights the canonical formal correspondence between the *rate of return* and the internal *string momentum*.

Here the tick variable h may be interpreted as a variable which extends along the extra dimension limited by the string size l_s . A natural consequence of the transform, Eq. (3), is the fulfillment of the boundary condition

$$P^{(1)}(\tau, 0) = 0, \quad (4)$$

which holds for any tick coordinate τ . Later on, we want to highlight effects of the rare events. For this purpose, we introduce a power-law q -deformed model

$$P_q^{(1)}(\tau, h) = f_q \left(\frac{p(\tau + h) - p(\tau)}{p(\tau + h)} \right), \quad h \in \langle 0, l_s \rangle \quad (5)$$

by means of the function

$$f_q(x) = \text{sign}(x) |x|^q, \quad q > 0. \quad (6)$$

The 1-end-point string has defined the origin, but it reflects the linear trend in $p(\cdot)$ at the scale l_s . Therefore, the 1-end-point string map $P_q^{(1)}(\cdot)$ may be understood as a q -deformed generalization of the *currency returns*. The illustration of the 1-end-point model is given in Fig. 1. The corresponding statistical characteristics displayed in Fig. 2 have been obtained on the basis of a statistical analysis discussed in Section 2.

Clearly, the situation with a long-term trend is partially corrected by fixing $P_q^{(2)}(\tau, h)$ at $h = l_s$. The open string with two end points is introduced via the nonlinear map which combines information about trends of p at two sequential segments

$$P_q^{(2)}(\tau, h) = f_q \left(\left(\frac{p(\tau + h) - p(\tau)}{p(\tau + h)} \right) \left(\frac{p(\tau + l_s) - p(\tau + h)}{p(\tau + l_s)} \right) \right), \quad h \in \langle 0, l_s \rangle. \quad (7)$$

The map is suggested to include boundary conditions of *Dirichlet type*

$$P_q^{(2)}(\tau, 0) = P_q^{(2)}(\tau, l_s) = 0, \quad \text{at all ticks } \tau. \quad (8)$$

In particular, the sign of $P_q^{(2)}(\tau, h)$ comprises information about the behavior differences of $p(\cdot)$ at three quotes (τ , $\tau + h$, $\tau + l_s$). The $P_q^{(2)}(\tau, h) < 0$ occurs for trends of the different sign, whereas $P_q^{(2)}(\tau, h) > 0$ indicates the match of the signs.

In addition to the variable $P_q^{(2)}(\tau, h)$ we introduced the conjugate variable $X_q^{(2)}(\tau)$ via the recurrent summation

$$X_q^{(2)}(\tau, h + 1) = X_q^{(2)}(\tau, h) + P_q^{(2)}(\tau, h - 1)[t(\tau + h) - t(\tau + h - 1)] \quad (9)$$

(here $t(\cdot)$ stands for a time-stamp corresponding to the quotation index τ in the argument). The above discrete form is suggested on the basis of the time-continuous Newton second law of motion $\dot{X}_q^{(2)}(t, h) = P_q^{(2)}(t, h)$ (written here for a unit mass). The form is equivalent to the imposing of the quadratic kinetic energy term $\frac{1}{2}(P_q^{(2)})^2$. Thus, the Hamiltonian picture [10] can be reconstructed in the following way;

$$\mathcal{H} = \frac{1}{2} \sum_{h=0}^{l_s} [(P_q^{(2)}(\tau, h))^2 - [\phi_{\text{ext}}(\tau, h + 1) - \phi_{\text{ext}}(\tau, h)] X_q^{(2)}(\tau, h)], \quad (10)$$

where $\phi_{\text{ext}}(\tau, h)$ is the external field term which depends on the transform of the currency rate [see e.g. Eq. (7)]. We pass from the continuum to discrete theory by means of the functional form

$$\dot{P}_q^{(2)} = - \frac{\delta \mathcal{H}}{\delta X_q^{(2)}(h)} = \phi_{\text{ext}}(\tau, h + 1) - \phi_{\text{ext}}(\tau, h) = P_q^{(2)}(\tau, h + 1) - P_q^{(2)}(\tau, h), \quad (11)$$

where $P_q^{(2)}(\tau, h)$ can be calibrated equal to $\phi_{\text{ext}}(\tau, h)$.

The discrete conjugate variable meets the *Neumann type* boundary conditions

$$X_q^{(2)}(\tau, 0) = X_q^{(2)}(\tau, 1), \quad X_q^{(2)}(\tau, l_s - 1) = X_q^{(2)}(\tau, l_s), \quad (12)$$

which is illustrated in Fig. 1(d).




Fig. 1. The illustrative examples of the currency data map for GBP/USD. The parts (a)–(d) constructed for date Fri, 31 Jul. 2009 time interval 15:06:37–15:43:09 GMT. Time evolution of symmetric ($P_{q=1}^{(1),S}$) and anti-symmetric ($P_{q=1}^{(1),A}$) component of the 1-end-point string of size $l_s = 1000$ calculated for $q = 1$ (by means of Eq. (18)). In (c), (d) we see the same data mapped by means of the partially closed 1-end-point string ($q = 1$) for $N_m = 10$, according to Eq. (26). (e) The calculation carried out for the 2-end-point string for $l_s = 1000$, $q = 6$ at some instant. We see that conjugate variable $X_{q=6}^{(2)}(\tau, h)$ satisfies the Neumann-type boundary conditions. (f) The instantaneous 2D-Brane state (date Fri, 31 Jul. 2009 15:11:47 GMT) is computed according to Eq. (23).

A more systematic way to obtain the 2-end-point string map represents the method of undetermined coefficients. The numerator of $q = 1$ can be chosen in the functional polynomial form of degree 2 with coefficients β_0, \dots, β_5 as follows:

$$P_{q=1, \text{Num}}^{(2)}(\tau, h) = \beta_0 p^2(\tau + h) + \beta_1 p^2(\tau) + \beta_2 p^2(\tau + l_s) \tag{13}$$

$$+ \beta_3 p(\tau)p(\tau + h) + \beta_4 p(\tau)p(\tau + l_s) + \beta_5 p(\tau + h)p(\tau + l_s). \tag{14}$$

Again, the Dirichlet conditions $P_{q=1, \text{Num}}^{(2)}(\tau, 0) = P_{q=1, \text{Num}}^{(2)}(\tau, l_s) = 0$ yield $P_{q=1, \text{Num}}^{(2)} = \beta_0(p(\tau) - p(\tau + h))(p(\tau + l_s) - p(\tau + h))$ with arbitrary β_0 . The overlooked denominator part of fraction $P_{q=1}^{(2)}$ then servers as a normalization factor.

Another interesting issue is the generalizing 1-end-point string to include the effect of many length scales

$$P_q^{(N_{l_s})}(\tau, h; \{l\}) = \prod_{i=1}^{N_{l_s}} f_q \left(\frac{p(\tau + l_i) - p(\tau + h)}{p(\tau + h)} \right), \tag{15}$$

which relies on the sequence $\{l\} \equiv \{l_i, i = 1, \dots, N_{l_s}\}$, including the end points ($\min_{i=1, \dots, N_{l_s}} l_i$ and $\max_{i=1, \dots, N_{l_s}} l_i$) as well as the $N_{l_s} - 2$ interior node points that divide the string map into the sequence of unfixed segments of non-uniform length (in general).

4. Symmetry with respect to the $p(\cdot) \rightarrow 1/p(\cdot)$ transform

The currency pairs can be separated into *direct* and *indirect* type. In a direct quote the *domestic* currency is the base currency, while the *foreign* currency is the quote currency. An *indirect* quote is just the opposite. Therefore, it would be interesting to take this symmetry into account. Hence, one can say that this two-fold division of the market network admits *duality symmetry*. Duality symmetries are some of the most interesting symmetries in physics. The term *duality* is used




Fig. 2. The variability in statistical characteristics caused by differences in topology and q . Calculated for the period Aug.–Sep., GBP/USD currency. The model with $q = 1$ has ability to reveal the currency long trend, on the other hand, the rare events are better visible for the 2-end-point string. The effect of the partial compactification with $N_m = 4$ [see Eq. (26)] is demonstrated in the third column (again for the 2-end-point string).

to refer to the relationship between two systems that have different descriptions but identical physics (identical trading operations).

Let us analyze the 1-end-point elementary string map when the currency changes from direct to indirect. The change can be formalized by means of the transformation

$$\hat{\mathcal{T}}_{id} : P\{p(\cdot)\} \rightarrow \bar{P}\{p(\cdot)\} \equiv P\{1/p(\cdot)\}, \tag{16}$$

For the 1-end-point map model of the string, Eq. (5), we obtained

$$\hat{\mathcal{T}}_{id} P_q^{(1)}(\tau, h) = \bar{P}_q^{(1)}(\tau, h) = f_q \left(\frac{p(\tau) - p(\tau + h)}{p(\tau)} \right). \tag{17}$$

Let us consider two-member space of maps $V_p^{(1)} = \{P_q^{(1)}, \bar{P}_q^{(1)}\}$. Importantly, we see that $\hat{\mathcal{T}}_{id}$ preserves the Dirichlet boundary conditions, in addition, the identity operator $\hat{\mathcal{T}}_{id}^2$ leaves the elements of $V_p^{(1)}$ unchanged. The space $V_p^{(1)}$ is closed under the left action of $\hat{\mathcal{T}}_{id}$. These ideas are straightforward transferable to the 2-end-point string points.

Now we omit the notation details and proceed according to Eq. (16). The map $P(\cdot)$ is decomposable into a sum of symmetric and antisymmetric parts

$$P^S = \frac{1}{2}(P + \bar{P}), \quad P^A = \frac{1}{2}(P - \bar{P}), \tag{18}$$

respectively. Due to of normalization by $1/2$, we get the projection properties

$$\hat{\mathcal{J}}_{id} P^S = P^S, \quad \hat{\mathcal{J}}_{id} P^A = -P^A. \tag{19}$$

To be more concrete, we choose $q = 1$ and obtain

$$P_{q=1}^{(1),S} = 1 - \frac{1}{2} \left[\frac{p(\tau)}{p(\tau+h)} + \frac{p(\tau+h)}{p(\tau)} \right], \quad P_{q=1}^{(1),A} = \frac{1}{2} \left[\frac{p(\tau)}{p(\tau+h)} - \frac{p(\tau+h)}{p(\tau)} \right] \tag{20}$$

and

$$P_1^{(2),A} = \frac{1}{2} \left[\frac{p(\tau)}{p(\tau+l_s)} - \frac{p(\tau+l_s)}{p(\tau)} + \frac{p(\tau+h)}{p(\tau)} - \frac{p(\tau)}{p(\tau+h)} + \frac{p(\tau+l_s)}{p(\tau+h)} - \frac{p(\tau+h)}{p(\tau+l_s)} \right], \tag{21}$$

$$P_1^{(2),S} = 1 + \frac{1}{2} \left[\frac{p(\tau+l_s)}{p(\tau)} + \frac{p(\tau)}{p(\tau+l_s)} - \frac{p(\tau)}{p(\tau+h)} - \frac{p(\tau+h)}{p(\tau)} - \frac{p(\tau+h)}{p(\tau+l_s)} - \frac{p(\tau+l_s)}{p(\tau+h)} \right].$$

We see that the $P_{q=1}^{(1),S}$ and $P_{q=1}^{(2),S}$ maps acquire formal signs of the systems with T -dual symmetry [2]. When the world described by the closed string of radius R is indistinguishable from the world of radius $\propto 1/R$ for any R , the symmetry manifests itself by $(R \pm \text{const.}/R)$ terms of the mass squared operator. The correspondence with our model becomes apparent when one assumes that R corresponds to the ratio $p(\tau)/p(\tau+h)$ in Eq. (20). However, we must also refer the reader to an apparently serious difference that in our model we do not consider for the moment, the compact dimension. One can also find in the option price dynamics some real examples of duality symmetry [11]. Concretely put–call duality which means “A call to buy foreign with domestic is equal to a put to sell domestic for foreign.” Also most questions will not spell out what is domestic or foreign but let you decide what is the underlying asset and which is the strike asset.

4.1. \mathcal{T}_{id} transform under the conditions of bid–ask spreads

Simply, the generalization can also be made with allowing for currency variables which appear as a consequence of the transaction costs [12]. The occurrence of ask–bid spread complicates the analysis in several ways. Instead of one price for each currency, the task requires the availability to two prices. The impact of ask–bid spread on the time-series properties has been studied within the elementary model [13].

Thus, for the purpose of a thorough and more realistic analysis of the market information, it seems straightforward to introduce generalized transform

$$\hat{\mathcal{J}}_{id}^{ab} P\{p_{ask}(\cdot), p_{bid}(\cdot)\} = \bar{P}\{1/p_{bid}(\cdot), 1/p_{ask}(\cdot)\}, \tag{22}$$

which converts to Eq. (17) in the limit of vanishing spread.

5. Mapping to the model of 2D-Brane

Clearly, there is a possibility to go beyond a string model towards more complex maps including alternative spread-adjusted currency returns. This is extension of string theory into the higher dimensions from the string lines into the membranes called D-Branes [14]. Formally, the generalized mapping onto the 2D-brane with the $(h_1, h_2) \in (0, l_s) \times (0, l_s)$ coordinates which vary along two extra dimensions could be proposed in the following form:

$$P_{2D,q}(\tau, h_1, h_2) = f_q \left(\left(\frac{p_{ask}(\tau+h_1) - p_{ask}(\tau)}{p_{ask}(\tau+h_1)} \right) \left(\frac{p_{ask}(\tau+l_s) - p_{ask}(\tau+h_1)}{p_{ask}(\tau+l_s)} \right) \right. \tag{23}$$

$$\left. \times \left(\frac{p_{bid}(\tau) - p_{bid}(\tau+h_2)}{p_{bid}(\tau)} \right) \left(\frac{p_{bid}(\tau+h_2) - p_{bid}(\tau+l_s)}{p_{bid}(\tau+h_2)} \right) \right).$$

The map constituted by the combination of “bid” and “ask” quotes is constructed to satisfy the Dirichlet boundary conditions

$$P_{2D,q}(\tau, h_1, 0) = P_{2D,q}(\tau, h_1, l_s) = P_{2D,q}(\tau, 0, h_2) = P_{2D,q}(\tau, l_s, h_2). \tag{24}$$

In addition, the above construction, Eq. (23), has been chosen as an explicit example, where the action of $\hat{\mathcal{J}}_{id}^{ab}$ becomes equivalent to the permutation of coordinates

$$\hat{\mathcal{J}}_{id}^{ab} P_{2D,q}(\tau, h_1, h_2) = P_{2D,q}(\tau, h_2, h_1). \tag{25}$$

Thus, the symmetry with respect to interchange of extra dimensions h_1, h_2 can be achieved through $P_{2D,q} + \hat{\mathcal{J}}_{id}^{ab} P_{2D,q}$. In a straightforward analogous manner one can get an antisymmetric combination $P_{2D,q} - \hat{\mathcal{J}}_{id}^{ab} P_{2D,q}$. For a certain instant of time we proposed illustration which is depicted in Fig. 1(b).

At the end of this subsection, we consider the next even simple example, where mixed boundary conditions take place. Now let the 2-end-point string be allowed to pass to the 1-end-point string by means of the homotopy $P_{q_1, q_2}^{(1,2)}(\tau, h, \eta) = (1 - \eta)P_{q_1}^{(1)}(\tau, h) + \eta P_{q_2}^{(2)}(\tau, h)$ driven by the parameter η which varies from 0 to 1. In fact, this model can be seen as a variant of the 2D-brane with extra dimensions h and η .

5.1. Partial compactification

In the frame of the string theory, the compactification attempts to ensure compatibility of the universe based on the four observable dimensions with twenty-six dimensions found in the theoretical model systems. From the standpoint of the problems considered here, the compactification may be viewed as an act of the information reduction of the original signal data, which makes the transformed signal periodic. Of course, it is not very favorable to close strings by the complete periodization of real input signals. Partial closure would be more interesting. This uses pre-mapping

$$\tilde{p}(\tau) = \frac{1}{N_m} \sum_{m=0}^{N_m-1} p(\tau + l_s m), \tag{26}$$

where the input of any open string (see e.g. Eq. (3), Eq. (7)) is made up partially compact.

Thus, data from the interval $\langle \tau, \tau + l_s(N_m - 1) \rangle$ are being pressed to occupy “little space” $h \in \langle 0, l_s \rangle$. We see that as N_m increases, the deviations of \tilde{p} from the periodic signal become less pronounced. The idea is illustrated in Fig. 1(c), (d). We see that the states are losing their original form (a), (b) are starting to create ripples.

For example, one might consider the construction of the $(\tilde{D} + 1)$ -brane

$$f_q \left(\frac{p(\tau + h_0) - p(\tau)}{p(\tau + h_0)} \right) \prod_{j=1}^{\tilde{D}} f_q \left(\frac{\tilde{p}_j^{(\pm)}(\tau + h_j) - \tilde{p}_j^{(\pm)}(\tau)}{\tilde{p}_j^{(\pm)}(\tau + h_j)} \right) \tag{27}$$

maintained by combining $(\tilde{D} + 1)$ 1-end-point strings, where partial compactification in \tilde{D} extra dimensions is supposed. Of course, the construction introduces auxiliary variables $\tilde{p}_j^{(\pm)}(\tau) = \sum_{m=0}^{N_m j-1} p(\tau \pm m l_{s,j})$.

6. Statistical investigation of 2-end-point strings

6.1. The midpoint information about string

In our present work, the strings and branes represent targets of physics-motivated maps which convert an originally dynamic range of currency data into the static frame. Of course, the data shaped by the string map have to be studied by statistical methods. However, the question remains open about the selection of the most promising types of maps from the point of view of interpretation of their statistical response.

Many of the preliminary numerical experiments we performed indicating that the 2-end-point strings with a sufficiently high q (in this work we focus on $q = 6$, but other unexplored values may also be of special interest) yield interesting statistical information including focus on rare events. Unfortunately, there is difficult or impossible to be exhaustive in this aspect. Fig. 3 shows how $\langle P_6^{(2)}(\tau, h) \rangle$ and the corresponding dispersion σ_{P_6} change with a string length.

6.2. The analysis of $P_q(l_s/2)$ distributions

The complex trade fluctuation data can be characterized by their respective statistical moments. In the case of the string map the moments of the ξ th order can be naturally considered at the half length

$$\mu_{q,\xi} = \langle |P_q^{(N)}(\tau, l_s/2)|^\xi / q \rangle. \tag{28}$$

The comparison of the results obtained for the 1-end point and 2-end point strings is depicted in Fig. 4. The remarkable difference in the amplitudes is caused by the manner of anchoring. The moments of longer strings are trivially larger.

6.3. Volatility vs. string amplitude

The volatility as described here refers to the standard deviation of currency returns of a financial instrument within a specific time horizon described by the length $l_s/2$. The return volatility at the time scale $l_s/2$ is defined by $\sigma_r(l_s/2) = \sqrt{r_2(l_s/2) - r_1^2(l_s/2)}$ using $r_m(l_s) = \sum_{h=1}^{l_s/2} [(p(\tau + h) - p(\tau + h - 1))/p(\tau + h)]^m$ for $m = 1, 2$. In Fig. 5, the rate of return




Fig. 3. The illustrative calculations carried out for EUR/USD currency. Figure shows the parts (a), (b) which include a view of two different epochs (and their different details). We see the variability of the mean statistical characteristics of the 2-end-point open string. The part (b) turns in sign, but remarkable exceptional scales corresponding to the local maxima and minima remain the same. The string length is expressed in real-time units calculated by means of Eq. (1). In part (c) we present anomalies – peaks roughly common for different currencies. These picture of anomalies are supplemented by dispersions of $P_q^{(2)}(l_s/2)$ (d).




Fig. 4. The mid-point fluctuations characterized by the statistical moments defined by Eq. (28). The calculations are carried out for GBP/USD currency rate, for the Aug.–Sep. period, for different kinds of strings for several lengths. We see that fluctuations become more significant as the string size increases. In addition, one may observe the 2-end-point string to be more suppressive to the fluctuations.

volatility computed at the scale $L = l_s/2$ demonstrates the linkage to the changes in the price trend represented by $P_6^{(2)}(l_s/2)$. Since the trend changes do not follow Gaussian distributions, we have used high q to analyze the impact of rare events. In Fig. 5, we show the identification of the semi-discrete levels of volatility by $q = 6$, while setting $q = 1$ does not uncover common attributes. In the future interest one can compare our return volatility computed from the string amplitude with some volatility estimators or GARCH type of models [15,16].

7. Intra-string statistical picture

The idea of the string related maps proposed here is the transformation of the original point object such as selected single price into a system of a prices from its admissible neighborhood. This changeover from a local to a non-local description directly extends econometrics belief that future prices are deducible from the price history of a given period. The results of the investigation of the intra-string statistics for a selected scale l_s (period) are presented in Fig. 6.

7.1. Time invariant strings and elementary statistics

To understand better the outputs of our numeric attempts, let us suppose to deal with short-time evolution of the currency $\ln p(\tau) = \ln p_0 + b\tau$ characterized by the linear-logarithmic parameter b . Surprisingly, after the substitution




Fig. 5. The scatterplot showing the relationship between the volatility $\sigma_\tau(\tau, l_s/2)$ and the string amplitudes $P_1^{(2)}(l_s/2)$ ($q = 1$) and $P_6^{(2)}(l_s/2)$ ($q = 6$), respectively. The separating effect at high q is visible. The plot indicates conservation or brake of the price trend $P_q^{(2)}(l_s/2)$ over the tick time $(\tau, \tau + l_s)$. We see that the trend becomes coupled with the occurrence of specific isolated values of the volatility calculated for $l_s = 10000$; period Aug.–Sep.

into the string, Eq. (7), the expressions collapse to the invariant (independent of τ) form

$$\begin{aligned}
 P_1^{(1)}(\tau, h) &= 1 - \exp(-hb), \\
 P_1^{(2)}(\tau, h) &= 1 - \exp[(h - l_s)b] - \exp(-hb) + \exp(-l_s b).
 \end{aligned}
 \tag{29}$$

It is quite interesting to look at the lowest order terms of Taylor series of this result around $b = 0$. Ignoring the terms of order b^4 or higher gives $P_1^{(1)} = bh - \frac{1}{2}b^2h^2 + \mathcal{O}(b^3)$, $P_1^{(2)} = b^2h(l_s - h) + \mathcal{O}(b^3)$. It demonstrates that the model of the 1-end-point string is more sensitive to the sign of b variations.

The elementary qualitative statistical model of the string can be obtained by taking the unexamined assumption that b fluctuates with the Gaussian probability density $(2\pi\sigma_b^2)^{-1} \exp[-b^2/(2\sigma_b^2)]$. The averaging of Eq. (29) with this weight yields

$$\langle P_1^{(1)} \rangle_{\text{Gauss}}(h) = 1 - \exp\left(-\frac{h^2\sigma_b^2}{2}\right),
 \tag{30}$$

$$\langle P_1^{(2)} \rangle_{\text{Gauss}}(h) = 1 - \exp\left(-\frac{h^2\sigma_b^2}{2}\right) - \exp\left(-\frac{(h - l_s)^2\sigma_b^2}{2}\right) + \exp\left(-\frac{l_s^2\sigma_b^2}{2}\right).
 \tag{31}$$

Eq. (30) predicts an increase in the h dependence as a consequence of the symmetric fluctuations in b . This finding does not agree with the result presented in Fig. 2, where the mean string output is biased by the mean trend.

7.2. Mapping of the periodic input signal

Simultaneously with giving numerical results obtained for statistical averages of data, it is instructive to briefly examine the string map of the signal of periodic form described by some elementary function. The input signal $p(\tau) = a_1 + a_2 \cos(\omega\tau)$ can be suitable for this purpose. Subsequently, the analytic calculation for the 2-end-point map can be carried out perturbatively under the requirement $a_2 \ll a_1$. The common form of average unifying formulas obtained for different integer q values can be written as

$$\langle P_q^{(2),(S)}(h) \rangle_{\text{cos}} = [\cos(h\omega) + \cos((h - l_s)\omega) - \cos(l_s\omega) - 1]^q \sum_{j=0}^{\infty} c_{q,j} \left(\frac{a_2}{a_1}\right)^{2(q+j)},
 \tag{32}$$




Fig. 6. The intra-string statistics for five currency pairs during the Aug.–Sep. period; the 2-end-point string is studied for $q = 6, l_s = 1000$. The mean $\langle P_6^{(2)}(l_s/2) \rangle$ (l.h.s. column) and dispersion (r.h.s. column) of $P_6^{(2)}(\tau, h)$. The specific scope brings analysis of the effects of symmetry and spread for statistics collected within constraint that GBP/USD base currency is trending up ($p_{bid}(0) \geq p_{ask}(0)$). The amplitude of $\langle P_6^{(2),A}(l_s/2) \rangle$ is very small (fluctuations are suppressed) as compared to the average of the symmetric part. On the other hand, $\langle P_6^{(2),S}(l_s/2) \rangle$ is not taken into the plot since it is indistinguishable from $\langle P_6^{(2)}(l_s/2) \rangle$.

where $c_{q,j}$ are the numerical coefficients which are not critical for further reasoning. The intuitive idea that $P_q^{(2),(A)}(\tau, h)$ discriminates fluctuations, which stems from the comparison of the symmetric and antisymmetric averages (see Fig. 6), is partially justified by the result $\langle P_q^{(2),(A)}(h) \rangle_{\cos} = 0$.

Interestingly, the calculation highlights the idea of the presence of the resonant lengths $l_s(n) = 2\pi n/\omega$ as $n = 1, 2, \dots$. This basic result motivated us to introduce the 2-end-point string model, that has the potential to identify characteristic dynamic scales represented here by $1/\omega$. It could be related with stochastic resonance for relevant currencies in the forex market. There is some similarity with stochastic resonance as a model for financial market crashes and bubbles found in Ref. [17]. It is important that anomalous aspect is absent in the statistical characteristics of the one-point strings.

7.3. String map in the representation of internal Fourier modes

Following the analysis of ultra-high-frequency financial data using advanced Fourier transformation as in Ref. [18] we assumed that each tick τ of the string may be represented by the sequence $P_q^{(2)}(\tau, h)$, $h = 0, 1, \dots, l_s - 1$ which can be transformed by means of the discrete Fourier transform

$$P_{\text{DFT},q}(k, \tau) = \sum_{h=0}^{l_s} P_q^{(2)}(\tau, h) \exp\left(-\frac{2\pi ikh}{l_s + 1}\right), \quad k = 0, 1, \dots, l_s. \tag{33}$$

Having done this, one can introduce the inverse transform

$$P_{\text{IDFT},q}(h, \tau) = \sum_{k=0}^{l_s} P_{\text{DFT},q}(\tau, k) \exp\left(\frac{2\pi ikh}{l_s + 1}\right), \tag{34}$$

which can be understood as a periodic extension of the input $P_q^{(2)}(\tau, h)$ with a period of the $(l_s + 1)$ ticks. Then $P_{\text{IDFT},q}(h, \tau)$ can be viewed as a portion of the original signal which curls up along the compact dimension of the closed string. Thus, the integer $h/(l_s + 1)$ has the meaning of a winding number of $P_{\text{IDFT},q}(h, \tau)$. The statistical mean values of $P_{\text{DFT},q}(k, \tau)$ calculated for different currencies are depicted in Fig. 7. The Fourier transform of the inherent string structure may serve to identify distinguishing features of currencies at selected time scale.

8. String polarized by the external field

In this section, we modify the two-point-string model, Eq. (7), in order to account for the transaction costs. This is some kind of analogy with a charged string which can be polarized by an external electric field [19]. A natural way is to consider the relation for the spread-adjusted return $(p_{\text{bid}}(\tau + h) - p_{\text{ask}}(\tau))/p(\tau + h)$ (written here for a long position). If we now routinely extend the 2-end-point, Eq. (7), we obtain

$$P_q^{\text{ab}}(\tau, h) = f_q \left(\left(\frac{p_{\text{bid}}(\tau + h) - p_{\text{ask}}(\tau)}{p(\tau + h)} \right) \left(\frac{p_{\text{bid}}(\tau + l_s) - p_{\text{ask}}(\tau + h)}{p(\tau + l_s)} \right) \right). \tag{35}$$

However this clearly violates, the Dirichlet boundary conditions, Eq. (8). The spread itself yields a negligible correction to the mean values.

The boundary conditions can be easily renewed by the subtraction $\tilde{P}_q^{\text{ab}}(\tau, h) = P_q^{\text{ab}}(\tau, h) - P_q^{\text{ab}}(\tau, 0)$. However, we show there exists a more fundamental alternative way which reflects a bid-ask difference and preserves the Dirichlet boundary conditions. The string states are polarized by the instant possibility to place a successful/unsuccessful buy order. For each h and $Y = A, S$ we construct the inequality constrained sequence

$$P_{(q,+)}^{(2),Y}(\tau + 1, h) = \begin{cases} P_q^{(2),Y}(\tau, h), & p_{\text{bid}}(\tau + l_s) \geq p_{\text{ask}}(\tau) \\ P_{(q,+)}^{(2),Y}(\tau, h), & \text{otherwise} \end{cases} \tag{36}$$

and non-buy contributions, respectively,

$$P_{(q,-)}^{(2),Y}(\tau + 1, h) = \begin{cases} P_q^{(2),Y}(\tau, h), & p_{\text{bid}}(\tau + l_s) < p_{\text{ask}}(\tau) \\ P_{(q,-)}^{(2),Y}(\tau, h), & \text{otherwise.} \end{cases} \tag{37}$$

In both cases it is supposed that $P_q^{(2),Y}(\tau, h)$ is calculated according an unconditioned model defined by Eq. (7). Now, to characterize the arbitrage opportunities, we introduced the statistical polarization measure in the form

$$g_{q,Y} = \left\langle \frac{\sum_{h=0}^{l_s} |P_{(q,+)}^{(2),Y}(\tau, h) - P_{(q,-)}^{(2),Y}(\tau, h)|}{\sum_{h=0}^{l_s} |P_{(q,+)}^{(2),Y}(\tau, h) + P_{(q,-)}^{(2),Y}(\tau, h)|} \right\rangle, \quad Y = A, S. \tag{38}$$




Fig. 7. The currency-specific slow ($k \leq 14$) intra-string averaged Fourier modes (their real and imaginary part separately) obtained by the temporal averaging over the Aug.–Sep. period. The mean real and imaginary parts calculated for five pairs, $q = 6$ and string length $l_s = 1000$.

The results of the extensive study of this measure are depicted in Fig. 8. These ideas could be extended to the case of market polarizations which are due to interactions amongst the agents in the market [20]. The agents could be characterized by the strings.

We continue the characterization of arbitrage opportunities by defining the (momentum) distance function between the strings as

$$d_q^{(Y)}(\tau) = \frac{1}{l_s + 1} \sum_{h=0}^{l_s} \left| P_{(q,+)}^{(Y)}(\tau, h) - P_{(q,-)}^{(Y)}(\tau, h) \right|. \tag{39}$$

In this case the statistics of string distances can be characterized by the customized variant of the well-known model of the correlation sum [7,21,22] or by the correlation structure patterns for identifying states of a financial market [23]. However, the motivation here differs from that given in these papers, where the intent was to analyze nonlinear relationships. The correlation sum shows the probability that the states of two strings or branes are localized within a certain distance. In our view we adjust the original formula to the string and brane models which, in addition, reflect the transactions involving profits. We define the measure

$$C_q^{(Y)}(\epsilon) = \frac{\langle \Theta(\epsilon - d_q^{(Y)}(\tau)) \rangle}{\int d\epsilon' \langle \Theta(\epsilon' - d_q^{(Y)}(\tau)) \rangle}, \tag{40}$$

where ϵ is the threshold distance, $\Theta(\cdot)$ is the Heaviside step function; here l_s plays the role of so called embedding dimension. The key concept surrounding this measure is the concept of the fractal dimension. In Fig. 9 we estimated a specific fractal dimension D_F as a slope of the dependence $d \ln C_q^{(Y)}(\epsilon) = D_F d \ln \epsilon$. However, as the figure shows, we have extended our




Fig. 8. The effect of string polarization on the mean characteristics calculated for selected currency pairs. One might identify a sudden increase of intra-string distinguishing features for the scales ranging from 3 to 5 h.




Fig. 9. The figure showing the statistics of inter-string distances. In part (a) the statistics collected and organized for the 2-end-point strings by means of the correlation sum (see Eq. (39)) for the currency pair GBP/USD and three time periods. As expected, the function $C_q^{(Y)}(\epsilon)$ monotonically decreases to zero as $\epsilon \rightarrow 0$. The index $D_F = 1.25 - 1.26$ varies over the periods. The values have been obtained by fitting over the estimated *universality range* ($10^{-7}, 2 \cdot 10^{-6}$) of ϵ . In part (b) the exponential dependences $C_{2D,1}(\epsilon)$ are obtained after the replacement of $d_q^{(Y)}(\tau)$ by $d_{2D,q}(\tau)$ in Eq. (40). We see that the inter-brane distances follow the exponential-type law with the highest variability across the short distances.

construction further and extend the concept of distance for 2D-branes as defined in Eq. (23). In a case like this we suggest generalization

$$d_{2D,q}(\tau) = \frac{1}{(l_s + 1)^2} \sum_{h_1=0}^{l_s} \sum_{h_2=0}^{l_s} |P_{2D,(q,+)}(\tau, h_1, h_2) - P_{2D,(q,-)}(\tau, h_1, h_2)|. \tag{41}$$

9. Inter-currency study: Map onto rotating strings

The incorporating of the mutual relations between the pairs into the mapping procedure represents a very challenging task. Let us study trading activity in the (I, J) plane, where I, J stands for indices of the currency pair described by two 2-end-point strings. The real time data are used instead of tick by tick (see Section 2) in order to maintain the consistency of prices quoted.

By continuing examples of the generalized distance concept (see Eq. (39)) and Eq. (41), we can introduce the inter-currency momentum distance function

$$d_{q,I,J}(t) = \frac{1}{l_s + 1} \sum_{h=0}^{l_s} |P_{q,I}^{(2)}(t, h) - P_{q,J}^{(2)}(t, h)|. \tag{42}$$

At higher dimensions, it is tempting to deal with angular momentum

$$M_{q,I,J}(t) = \sum_{h=0}^{l_s} [P_{q,I}^{(2)}(t, h)X_{q,J}^{(2)}(t, h) - P_{q,J}^{(2)}(t, h)X_{q,I}^{(2)}(t, h)]. \tag{43}$$

The momentum calculation can be interpreted as a measure of the rotational information flows between the currency pairs. In Fig. 10 we present the results achieved for two alternatives $(I, J) = (EUR/USD, GBP/USD)$, and $(I, J) = (GBP/USD, USD/JPY)$. We see that calculation of $D_{q,I,J}$ and $M_{q,I,J}$ yields oscillatory (i.e. seasonal) behavior. Simultaneously, the concept of distance and moment has been extended to analyze the impact of spread. Analogously, as in the previous cases, the distance between the ask and bid strings may be defined

$$d_{q,I}^{ab}(t) = \frac{1}{l_s + 1} \sum_{h=0}^l |P_{q,ask}^{(2)}(t, h) - P_{q,bid}^{(2)}(t, h)|, \tag{44}$$




Fig. 10. The inter-string distance and angular momentum (see Eqs. (42) and (43)) for specified currency pairs. The long term outlook compared with a detailed one-week view supplemented by the results obtained for spread according to Eqs. (44) and (45). The results are compared with the Gâteaux derivative (see Eq. (50), where p corresponds to EUR/USD and ψ to GBP/USD pair), and the derivative is evaluated at $h = l_s/2$. The calculation is carried for the string of the 1 h time length and investigation of the Aug.–Sep. period.

$$M_{q,I}^{ab}(t) = \sum_{h=0}^{l_s} \left[P_{q,ask}^{(2)}(t, h) X_{q,bid}^{(2)}(t, h) - P_{q,bid}^{(2)}(t, h) X_{q,ask}^{(2)}(t, h) \right]. \tag{45}$$

Here

$$P_{q,ask}^{(2)} \equiv P_q^{(2)}|_{p \rightarrow p_{ask}}, \quad P_{q,bid}^{(2)} \equiv P_q^{(2)}|_{p \rightarrow p_{bid}} \tag{46}$$

are obtained by substituting expressions above in Eq. (7). With the help of Eq. (9) and $P_{q,ask}^{(2)}, P_{q,bid}^{(2)}$ we construct iteratively $X_{q,ask}^{(2)}$ and $X_{q,bid}^{(2)}$. As one can see from Fig. 10, the differences measured in terms of $M_{q,I,J}(t)$ are very subtle. There is evidence of intercoupling of the spread and currency dynamics. The fundamental role in the string theory is played by the *Regge slope parameter* α' (or inverse string tension). This has a proper analogy with our approach where we introduced a slope in terms of the angular momentum

$$\alpha'_{q,I,J} = \frac{\langle |M_{q,I,J}| \rangle}{2\pi l_s^2}. \tag{47}$$

For the $l_s = 1$ h string pair we obtained $\alpha'_{6,\text{EUR/USD,GBP/USD}} = 5.07 \cdot 10^{-54} (2\pi)^{-1} \text{h}^{-2}$, $\alpha'_{6,\text{GBP/USD,USD/JPY}} = 1.55 \cdot 10^{-53} (2\pi)^{-1} \text{h}^{-2}$, $\alpha'_{6,\text{USD/JPY,EUR/USD}} = 1.34 \cdot 10^{-53} (2\pi)^{-1} \text{h}^{-2}$ much larger than the spread $\alpha'_{6,\text{GBP/USD}} = 1.16 \cdot 10^{-55} (2\pi)^{-1} \text{h}^{-2}$. However, it is worth noting that relation, Eq. (47), should be understood as an estimate since there is no statistical mean of the type $\langle |M_{\dots}| \rangle$ in the original specification. The problem of estimation of the *slope parameter* arises from the fact that in the original model nonaveraged angular momentum is divided by the square of the mass instead of l_s^2 . Herein, we have no idea how to measure the mass of the string, or how to verify the fact that the string mass is proportional to l_s .

10. Differentials of string map

Gâteaux derivative is a generalization of the concept of a directional derivative in the differential calculus. In our study the concept can be viewed as a systematic way in the generation of more structured maps expressing more information about the structure of data we deal with. Given the string map $P(\cdot)$, the m -th Gâteaux derivative of $P(\cdot)$ in the “direction” of $\psi(\cdot)$ (unspecified yet series) is defined as follows:

$$d^m P(\{p\}; \{\psi\})(\tau, h) = \frac{d^m}{d\epsilon^m} P(\{p(\tau, h) + \epsilon \psi(\tau, h)\})|_{\epsilon \rightarrow 0}. \quad (48)$$

For $q = 1$ the calculation gives

$$dP_1^{(1)}(\{p\}; \{\psi\})(\tau, h) = \frac{1}{p(\tau + h)} \left[\frac{p(\tau)\psi(\tau + h)}{p(\tau + h)} - \psi(\tau) \right] \quad (49)$$

and

$$dP_1^{(2)}(\{p\}; \{\psi\})(\tau, h) = \psi(\tau) \left(\frac{1}{p(\tau + l_s)} - \frac{1}{p(\tau + h)} \right) + \psi(\tau + h) \left(\frac{p(\tau)}{p^2(\tau + h)} - \frac{1}{p(\tau + l_s)} \right) + \frac{\psi(\tau + l_s)}{p^2(\tau + l_s)} (p(\tau + h) - p(\tau)). \quad (50)$$

By going to the second order we obtained

$$d^2 P_1^{(2)}(\{p\}; \{\psi\})(\tau, h) = \frac{2\psi(\tau + h)}{p^2(\tau + h)} \left[\psi(\tau) - \frac{p(\tau)\psi(\tau + h)}{p(\tau + h)} \right] \quad (51)$$

and

$$d^2 P_1^{(2)}(\{p\}; \{\psi\})(\tau, h) = \frac{2\psi(\tau + l_s)}{p^2(\tau + l_s)} [\psi(\tau + h) - \psi(\tau)] + \frac{2\psi(\tau + h)}{p^2(\tau + h)} \left[\psi(\tau) - \frac{p(\tau)\psi(\tau + h)}{p(\tau + h)} \right] + \frac{2\psi^2(\tau + l_s)}{p^3(\tau + l_s)} [p(\tau) - p(\tau + h)]. \quad (52)$$

Surprisingly, the generalized differentiation generates maps which satisfy the Dirichlet boundary conditions

$$d^m P_1^{(1)}(\{p\}; \{\psi\})(\tau, 0) = 0, \quad m = 1, 2; \quad (53)$$

$$d^m P_1^{(2)}(\{p\}; \{\psi\})(\tau, 0) = d^m P_1^{(2)}(\{p\}; \{\psi\})(\tau, l_s) = 0. \quad (54)$$

Many alternative ways exist to exploit the models with the auxiliary field $\psi(\cdot)$. The field can be related to, e.g., (i) models which place emphasis on the currency margins determined by some adaption process; (ii) on the spread in a style of Section 8 with $\psi(\tau) = p_{\text{bid}}(\tau) - p_{\text{ask}}(\tau - l_s)$. (iii) The benchmark setting represents $\psi = 1$, (iv) the periodic function $\psi(\tau)$ can model the action of the compact. In this intuition supporting case, one can see that the generalized derivative modifies the original map as follows:

$$dP_1^{(1)}(\{p\}; \{\psi\})(\tau, h)|_{\psi=1} = -\frac{P_1^{(1)}(\tau, h)}{p(\tau + h)}, \quad (55)$$

$$dP_1^{(2)}(\{p\}; \{\psi\})(\tau, h)|_{\psi=1} = \left(\frac{1}{p(\tau + h)} + \frac{1}{p(\tau + l_s)} \right) P_1^{(2)}(\tau, h). \quad (56)$$

In Fig. 10, we present the idea, where p, ψ are represented by two currencies, their mutual influence is studied within the first-order differential model described by Eq. (50).

11. Conclusions

We shown that the string theory may motivate the adoption of the nonlinear techniques of the data analysis with a minimum impact of justification parameters. The numerical study recovered interesting fundamental statistical properties of the maps from the data onto string-like objects. The remarkable deviations from the features known under the notion of the efficiently organized market have been observed, namely, for high values of the deformation parameter q .

The main point here is that the string map gives a geometric interpretation of the information value of the data. The model of the string allows one to manipulate the information stored along several extra dimensions. We started from the theory of the 1-end-point and 2-end-point string, where we distinguished between the symmetric and antisymmetric variants of the maps. In this context, it should be emphasized that duality is a peculiar property of the suggested maps, not data alone. On the contrary with call–put duality in option pricing there could be some relation with string duality. The numerical analysis of the intra-string statistics was supplied qualitatively by the toy models of the maps of the exponential and periodic data inputs. Most of the numerical investigations have been obtained for the open topology; however, we described briefly the ways to partial compactification. The data structures can also be mapped by means of the curled dimension which arises as a sum of periodic data contributions. The idea of the compactified strings can be realized as well by the application of the inverse Fourier transform of the original signal. The interesting and also challenging task represents finding of link between string map and log-periodic behavior of speculative bubbles of the stock market indices [24,25]. It would be also interesting to examine R/S analysis of the Hurst exponents [26,27] for the case of finite strings instead of the usual point prices.

The study of string averages exhibited occurrences of the anomalies at the time scales proportional to the string length. We showed that global and common market timescales can be extracted by looking at the changes in the currencies. The extensions of the string models of branes including ask/bid spread were discussed. The membrane 2d-brane approach could be also helpful e.g. for computations of the volatility surface in option pricing [28]. We studied the relationship between the arbitrage opportunities and string statistics. We showed that extraction of the valuable information about the arbitrage opportunities on given currency could be studied by means of the correlation sum which reflected the details of the occupancy of phase–space by differently polarized strings and branes. In addition, we presented several physics and geometry motivated methods of analysis of the coupling between the currency pairs. The results led us to believe that our ideas and methodology can contribute to the solution of the problem of the robust portfolio selection. As we have seen, the complex multi-string structures produced by the generalized derivatives of strings cannot be easily grasped by the intuitive principles. We believe, the method affords potential to be used in the practical applications, where arbitrage selection bias should be taken into account. In the future research we will focus on the real application of our string approach. The main goal will be finding fundamental principles of the dynamics and following prediction in behavior to the financial forex market.

12. List of terms

- String theory: is an active research framework in particle physics where particles are rather 1-dimensional oscillating open or closed lines “strings”.
- Brane theory: are membranes of different dimensionality from a one-dimensional membrane which is in fact a string lines, including 2, 3 or more dimensional membranes.
- Extra dimension: string theory predicts extra dimensions, in classical string theory the number of dimensions is not fixed by any consistency criterion.
- Conjugate variable: are pairs of variables mathematically defined in such a way that they become Fourier transform duals of one-another.
- T-duality: is a symmetry of quantum field theories with differing classical descriptions, of which the relationship between small and large distances in various string theories is a special case.
- Compact dimension: is curled up in itself in very small Planck length and the fact that the dimension is smaller than the smallest particle means that it cannot be observed by conventional means.
- Regge slope parameter: was introduced in the quantum theory of string, and its relation to the string tension involves \hbar .
- Gâteaux derivative or directional derivative is often used to formalize the functional derivative commonly used in the calculus of variations and physics.

Acknowledgments

All calculations were done on the computational cluster designed by the company FURT Solutions, s.r.o., Kosice, Slovak Republic, <http://www.furt.sk>. For more information about the progress of our new string application work to financial markets please follow the line <http://www.sors.com/people/pincak/>.

References

- [1] D. McMahon, *String Theory Demystified*, The McGraw-Hill Companies, Inc., 2009.
- [2] B. Zwiebach, *A First Course in String Theory*, Cambridge University Press, 2009.
- [3] J. He, M. Deem, *Physical Review Letters* 198701 (2010) 188.

- [4] W.S. Jung, O. Kwon, F. Wang, T. Kaizoji, H.T. Moon, H.E. Stanley, *Physica A* 387 (2008) 537.
- [5] M. Eryiğit, R. Eryiğit, *Physica A* 388 (2009) 3551.
- [6] P.R. Krishnaiah, L. Kanal, *Classification, Pattern Recognition, and Reduction of Dimensionality*, North-Holland, Amsterdam, 1982.
- [7] P. Grassberger, I. Procaccia, *Physica D* 9 (1983) 189.
- [8] Y. Ding, X. Yang, A.J. Kavs, J. Li, *International Journal of Trade, Economics and Finance* 1 (2010) 320.
- [9] J. Polchinski, *String Theory*, Cambridge University Press, 1998.
- [10] M.B. Green, J.H. Schwarz, E. Witten, *Superstring Theory*, Cambridge University Press, 1987.
- [11] J. Fajardo, E. Mordecki, *Quantitative Finance* 6 (2006) 219.
- [12] W.H. Wagner, M. Edwards, *Financial Analysts Journal* 49 (1993) 65.
- [13] R. Roll, *The Journal of Finance* 39 (1984) 1127.
- [14] C.V. Johnson, *D-Branes*, Cambridge University Press, 2003.
- [15] B. Bollen, B. Inder, *Journal of Empirical Finance* 9 (2002) 551.
- [16] P. Molnar, *International Review of Financial Analysis* (in press) Corrected Proof, Jul 2011 <http://dx.doi.org/10.1016/j.irfa.2011.06.012>.
- [17] A. Krawiecki, J.A. Holyst, *Physica A* 317 (2003) 597.
- [18] I. Giampaoli, W.L. Ng, N. Constantinou, *Finance Research Letters* 6 (2009) 47.
- [19] P.O. Kazinski, *Journal of Experimental and Theoretical Physics* 101 (2005) 270.
- [20] S. Sinha, S. Raghavendra, *Advances in Artificial Economics* 584 (2006) 177.
- [21] J.Y. Campbell, A.W. Lo, A.C. MacKinlay, *The Econometrics of Financial Markets*, Princeton University press, 1997.
- [22] N. Bigdeli, K. Afshar, *Physica A* 388 (2009) 1577.
- [23] M.C. Münnix, T. Shimada, R. Schäfer, F.L. Thomas, H. Seligman, T. Guhr, H.E. Stanley, [arXiv:1202.1623v1](https://arxiv.org/abs/1202.1623v1), 8 February 2012.
- [24] W.X. Zhou, D. Sornette, *Physica A* 330 (2003) 543.
- [25] W.X. Zhou, D. Sornette, *Physica A* 337 (2004) 243.
- [26] B. Mandelbrot, *Review of Economics and Statistics* 53 (1971) 225.
- [27] M.A.S. Granero, J.E.T. Segovia, J.G. Pérez, *Physica A* 387 (2008) 5543.
- [28] M.J. Kim, S.Y. Lee, D.I. Hwang, S.Y. Kim, I.K. Ko, *Physica A*, 389 14 (2762).